

AAA Foundation for Traffic Safety

FACT SHEET

Prevalence of Marijuana Involvement in Fatal Crashes: Washington 2010- 2014

Background

- In November 2012, the citizens of the state of Washington approved by popular vote ballot Initiative 502, which allows adults aged 21 years and older to possess up to 1 ounce of marijuana for personal use
 - The new law became effective on December 6, 2012
- Not much is known about the prevalence of driving after using marijuana or the prevalence of recent marijuana use among drivers involved in crashes primarily due to data limitations. In January 2016, the Washington Traffic Safety Commission (WTSC) made available the results of toxicology tests for THC performed on drivers involved in fatal crashes in the state of Washington, and appended these new data to their database of all drivers involved in fatal crashes statewide
 - This study uses these new data from the WTSC to estimate the proportion of drivers in fatal crashes in the state of Washington who had a detectable concentration of THC in their blood at the time of the crash and to investigate whether that proportion changed after Washington Initiative 502 took effect on December 6, 2012

Objective

- The purpose of this study was to quantify the prevalence of marijuana involvement in fatal crashes in the state of Washington in years 2010 – 2014 and to investigate whether the prevalence changed after Washington Initiative 502, which legalized recreational use of marijuana for adults aged 21 years and older, took effect on December 6, 2012

Methods

- The data examined were obtained from the WTSC and comprised a census of all motor vehicle crashes that occurred on public roads in the state of Washington and resulted in a death within 30 days
- Examined the presence and concentration of delta-9-tetrahydrocannabinol (hereafter *THC*), the main psychoactive chemical in marijuana, in the blood toxicological test results of drivers involved in fatal crashes
- THC presence and concentration in the subset of drivers whose blood was not tested or whose test results were unavailable were estimated using the method of multiple imputation. The imputation method explicitly accounted for changes implemented during the study period in the cutoff levels used in the state laboratory for detection of THC




Saving lives
through research
and education



Key Findings

- Statewide, 3,031 drivers were involved in fatal crashes in years 2010 – 2014
- Overall, considering both the actual blood toxicology test results and imputed results, an estimated 303 drivers—**10.0%** of all drivers involved in fatal crashes in Washington between 2010 and 2014—had detectable THC in their blood at the time of the crash
- Of all THC-positive drivers involved in fatal crashes, an estimated 34.0% had neither alcohol nor other drugs in their blood, 39.0% had detectable alcohol in addition to THC, 16.5% had other drugs in addition to THC, and 10.5% had had both alcohol and other drugs in addition to THC in their blood at the time of the crash
- From 2010 through 2013, the estimated number and proportion of drivers involved in fatal crashes who had a detectable concentration of THC in their blood ranged from a low of 48 (7.9%) to a high of 53 (8.5%)
 - The number and proportion both doubled from 49 (8.3%) in 2013 to 106 (17.0%) in 2014
- Analysis of trends over time before and after Initiative 502 took effect indicate that the proportion of drivers positive for THC was generally flat before Initiative 502, but began increasing significantly approximately 9 months after the effective date of Initiative 502

Figure 1. Quarterly average proportion of drivers involved in fatal crashes who were positive for THC and modeled seasonally-adjusted linear trend before and after Washington Initiative 502 took effect on 6 December 2012 legalizing recreational use of marijuana for adults aged 21 years and older, Washington, 2010 – 2014



Data: Washington Traffic Safety Commission, 2010 – 2014.

Drivers positive for THC based on results of blood toxicological tests. Results imputed 10 times when driver was not tested or test results were unknown; results reflect averages from 10 imputed values for each driver. Model-based predictions are from binomial regression model with identity link function, indicator variables for seasons, and a two-part linear spline with change in slope on 5 September 2013 (39 weeks after effective date of Initiative 502).

For more information on this study and the AAA Foundation's other traffic safety research and materials, please visit AAAFoundation.org.

May 2016

